


---




ESE

## 8. Experiment Planning




# Planning a Controlled Experiment

- Hypothesis formulation
- Variables selection
  - Independent variables
    - Factors
      - Treatments, Parameters
    - Noises
  - Dependent Variables
- Experiment design
  - Randomization
  - Blocking
  - Balancing
  - Standard design types 22
- Instrumentation
- Threats to validity evaluation



# Context selection

- ❑ Offline vs. online
- ❑ Student vs. professionals
- ❑ Toy vs. real problems
- ❑ Specific vs. general



# Hypotheses formulation

## Null hypothesis


$H_0$  : There is no underlining trend or pattern in the experiment setting.

*Usually,  $H_0$  is the statement that experimenters want to reject.*

*E.g.,  $H_0: \mu_1 == \mu_2$*

*“==” is preferred, because it is easier to evaluate than other relation operators.*

**Serious scientists should make all acts to prove that  $H_0$  is not to reject.**



# Hypotheses formulation

## Alternative hypothesis

$H_1$  : There is a significant trend or pattern in the experiment setting.


*Usually, the statement in favor of which  $H_0$  is rejected.*

*E.g.:*

1)  $H_1: \mu_1 \neq \mu_2$

2)  $H_1: \mu_1 < \mu_2$  ←

***Often but not necessarily, 2) is the  $H_1$  in industry.***



# Hypotheses formulation:


## Note

$$!H_0 !\leftrightarrow H_1$$

If, with respect to data from a given experiment, you cannot prove that  $H_0$  is reasonable true, this does not mean that  $H_1$  is reasonable true.

Recall that we do not work in an axiomatic system where  $!(=) \leftrightarrow <>$ .

See a next chapter on What is an hypothesis, hypotheses testing, and related risks.




# Hypotheses formulation: Example

$$H_{0EC} : \mu_{1EC} = \mu_{2EC}$$

$$H_{1EC} : \mu_{1EC} \neq \mu_{2EC}$$

$H_0$ (Efficiency): CR and FTI perform  
**insignificantly different** *in detecting defects.*

$H_1$ (Efficiency): CR and FTI perform  
**significantly different** in detecting defects.



# Hypotheses formulation: Example


$$H0_{ES} : \mu1_{ES} = \mu2_{ES}$$

$$H1_{ES} : \mu1_{ES} \neq \mu2_{ES}$$

H0(Effectiveness): CR and FTI perform **insignificantly different** in detecting defects.

H1(Effectiveness): CR and FTI perform **significantly different** in detecting defects.





# Hypotheses formulation: Refinement

1)  $H_1: \mu_1 \neq \mu_2$

2)  $H_1: \mu_1 < \mu_2$

3)  $H_1: \mu_1 \leq 1,2 * \mu_2$

4)  $H_1: \mu_1 \leq 1,2 * \mu_2 + 0,5$

5)  $H_1: \mu_1 \leq 1,2 * \mu_2^{0,1}$

...



# Variables selection <sup>(1/2)</sup>


Dependent and independent variables must be chosen **before** designing the experiment.

- ✓ Choice of independent variables
- ✓ Choice of dependent variables



**Usually, dependent variables come first (from Goals)**


Once we have chosen those variables, and levels and treatments, experiment strategy and design can be defined.



# Variables selection

Usually we cannot afford all the input variables and all their treatments in one time. In order to manage the experiment complexity, we proceed by step-wise-refinement:

- ✓ **Identify and verify independent variables**
  - ✓ (1<sup>st</sup> experiment).
- ✓ **Reduce the number of significant variables**
  - ✓ Choose design factor[s] as that [those] input variable[s] that more than other ones affect[s] outcomes; choose parameters, and identify disturbs and noises.
- ✓ **Reduce the number of treatments**
  - ✓ For each design factor choose (“*Constant effects model*”), or “*Fixed effects model*”, or select at random (“*Random effects model*” or “*Variance components model*”) a few (e.g. 2) of its alternatives.
- ✓ **Reduce the number of blocks**
  - ✓ From complete to incomplete “blocked” “factorial” design.




# Subject Selection

Generally we select a sample of  $n$  subjects (“**sample size**”) from a population.

In order to generalize the results to the desired population, the selection must be representative for that population.


The population should be studied carefully before taking a sample.



# Defining the Population


We define our population:

- **Experience-based**: Based on some given materials in advance, or some certain inputs.
- **Environment-based**: Based on being affected by a certain environment and conditions.



# Subject Selection as Probability Sample

If the probability of selecting each subject is known then the selection is a **probability sample**.



# Probability Samples


**Simple random sampling:**  $n$  subjects are selected from a list of the population at random.

**Systematic sampling:** The first subject is selected from the list at random, hence the following  $n-1$  ones are selected.

**Convenience sampling:** The nearest and most convenient people are selected as subjects.

**Quota sampling:** This type of sampling is used to get subjects from various elements of a population. Hence, Convenient sampling is normally used for each element.

**Stratified random sampling:** The population is divided in a number of groups or strata with a known distribution between the groups. Random selection is then applied within the strata.



# Experimental Design: Basic Principia

1. Replication
2. Randomization
3. Blocking
4. Balancing





# Replication

Replication (*of the basic experiment*) means to repeat the basic experiment, i.e. the minimum set of the elementary experiments. [See also *Replication of an experiment*.]

Replication allows us:

- 1) to estimate the *experimental error* (hence the **significance** of a result),
- 2) to estimate more precisely the *impact of a factor on the output*. e.g., by diminishing the **variance** of the **sample mean** with respect to the variance of the **single observation**.

---

(See impact on errors in Ch. 10)




# Randomization

**Randomization** is the mile stone that allows using **statistics** in experimentation (errors are assumed independent and random variables).


- It allows to average on factors that may otherwise show their presence.
- It is used in allocating objects, subjects, and test order.

*When complete randomization is not possible to enact, we have to adopt specific planning statistic methods.*



# Blocks


- They allow to improve the precision from comparison between *interesting* factors.
- They also concern **disturbing factors**, *known and controllable factors* that probably have an effect, but we are not interested in that effect.
  - ❖ We block with respect to such a factor, when we arrange the experiment in a way that **in a block that factor is constant**.
  - ❖ *We are not expected to study effects between blocks.*



# Balancing

Balancing subjects in an experiment means to assign the treatments so that each treatment has the same number of subjects.

In this case, the statistical analysis is simplified and strengthened.



# Usage of Statistical vs. Non-statistical Techniques

- Use also **non-statistical knowledge** about the problem.
- Adopt experimental plans and data analysis as **simple** as possible.
- Distinguish between practical and statistical **significance**.
- In order to improve knowledge, experiments are **iteratively** realized.

# Experiment Design Types

**One factor** //(n=>2 treatments)

- **Randomized [Incomplete Block] Design**

// One treatment per subject

- **Simple** [IB] Randomized Design, S[IB]RD

// One treatment per object

*Objects are assigned randomly to treatments*

- **Completely** [IB] Randomized Design, C[IB]RD

// All treatments are applied to an object (\*)

*Subjects are assigned randomly to treatments.*

- **Randomized Complete Block Design**

// Each subject uses all the treatments (\*)


- **[Simple] RCBD** // One treatment is applied to an object.

- **[Completely] RCBD** // All treatments are applied to an object. (\*)

(\*) *The total order in which subjects use the treatments is assigned at random.*

Treatments vs. Subjects


Treatments vs. Objects



# Experiment Design Types

## More than one factor

- Two treatments per factor
  - Two factors
    - $2^2$  factorial design
    - Two-stage nested factorial design
  - More than two factors (k)
    - $2^k$  factorial design
    - $2^k$  fractional factorial design.
- More than 2 treatments per factor (n)




# Undesired Variables

It is necessary to evaluate the presence of undesired variables and make decision about how the handle them.

- For instance: Are there software professionals and non-professionals among the student subjects participating to an academic software engineering experiment? How may level of experience can you define? How to handle each of them?






# Experiment Design Types

CONDITIONS		DESIGN
1 factor of interest - 2 treatments - $n$ treatments	All other project parameters can be fixed	- Simple randomized exp. - Paired comparison <i>Completely randomized design</i>
	<i>There is variability between subjects</i>	<i>Randomized complete block design</i>
K factors of interest (2 or $n$ treatments)	There are undesired variations  $n^k$ experiments	Blocked factorial design  - Factorial design - Nested design
	There are desired variations only  $= [ < ] n^k$ experiments	[Fractional] Factorial design

TO UPDATE



# Instrumentation and experiment materials

Is concerned with:

- ❑ the arrangement of **objects**, and **forms** and **guidelines** to give to experiment subjects
- ❑ **measurement tools** and further **supports** for *data collection*.


E.g. :

- ❑ Selecting program-code to read or test for defect detection; seeding the code with defects.
- ❑ Arranging forms for defining and identifying defects found.
- ❑ Arranging the net to collect the submission times.

---




# Planning the training of subjects



# Types of Validity

- **Internal validity**: wants to make sure that a statistical relationship between inputs and outcomes is a **causal relationship**.
- **External validity**: concerns the *generalization* of the results *outside* the scope of the study.
- **Construct validity**: is concerned with relationship between the level of the *theory* and the level of the *observation*; depends on the adequacy of used *measurement* models.
- **Conclusion validity**: It is concerned with relationship between treatment and outcome.



# Evaluating the Validity Threats

(that derive from the planned choices)

[See 8.1](#)

[See 8.2 for further details](#)